

What to do

Advice on childhood illnesses



Go to school; if needed, get treatment as required



Some restrictions for school attendance



Don't go to school and see the GP

What it's called?	What it's like	Going to school	Getting treatment	More advice
Asthma flare up	Worsening of usual symptoms		Follow your care plan	Stay off school if feeling too unwell
Chicken pox	Rash begins as small, red, flat spots that develop into itchy fluid-filled blisters		Pharmacy	Back to school five days after on-set of the rash, or when all lesions are crusted over
Common cold	Runny nose, sneezing, sore throat		Pharmacy	Ensure good hand hygiene
Conjunctivitis	Teary, red, itchy, painful eye(s)		Pharmacy	Try not to touch eye to avoid spreading*
Flu	Fever, cough, sneezing, runny nose, headache body aches and pain, exhaustion, sore throat		Pharmacy	Ensure good hand hygiene*
German measles/ Rubella	Fever, tiredness. Raised, red, rash that starts on the face and spreads downwards		GP	Back to school four days after the rash has started, if feeling well enough. Please let the school know, as pregnant members of staff may be affected.
Glandular fever	High temperature, swollen glands, sore throat; usually more painful than any before		GP	*
Hand, foot & mouth disease	Fever, sore throat, headache, small painful blisters inside the mouth on tongue and gums (may appear on hands and feet)		GP	Attend school while monitoring symptoms
Head lice	Itchy scalp (may be worse at night)		Pharmacy	
Impetigo	Clusters of red bumps or blisters surrounded by area of redness		GP	Back to school when lesions crust or 48 hours after start of antibiotics
Measles	Fever, cough, runny nose, watery inflamed eyes, rash		GP	Back to school four days from on-set of rash, and if feeling well enough
Ringworm	Red ring shaped rash, may be itchy, rash may be dry and scaly or wet and crusty		Pharmacy	
Scabies	Itching rash, commonly between fingers, wrists, elbows, arm		GP	Back to school after first treatment
Scarlet fever	Fever, unwell, red tongue, sandpaper rash		GP	Exclusion period during first 24 hours of antibiotics*
Shingles	Pain, itching, or tingling along the affected nerve pathway. Blister-type rash		GP	Only stay off school if rash is weeping and cannot be covered
Sickness bug/ diarrhoea	Stomach cramps, nausea, vomiting and diarrhoea		Pharmacy	Contact GP if symptoms persist after 48 hours
Threadworms	Intense itchiness around anus		Pharmacy	Ensure good hand hygiene
Tonsillitis/ Strep throat	Intense sore throat for more than 1 day		Pharmacy	See GP if temperature lasts more than 48 hours or cannot swallow . Only need to stay off school if feeling too unwell
Whooping cough	Violent coughing, with a 'whoop'		GP	Exclusion period during first five days of antibiotics
Covid				Visit www.nhs.uk for latest guidance

*Attend school while monitoring symptoms.

This information is a guide and has been checked by health professionals. If you are unsure about your child's wellbeing, we recommend you contact your pharmacy or GP to check.

Should my child go to school/nursery today?

Hertfordshire and West Essex Healthier Together for further information

<https://www.hwehealthiertogether.nhs.uk/parentscarers>

Working in partnership with

